

Tilmicosin and Monensin Medicated Cattle Feed – BRD and Coccidiosis

Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS

For prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii* and control of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni* in groups of cattle fed in confinement for slaughter, where active BRD has been diagnosed in at least 10% of the animals in the group.

ACTIVE DRUG INGREDIENTS

Tilmicosin (as tilmicosin phosphate) ^c	568 to 757 grams per ton ^{*, a}
Monensin, USP ^d	10 to 40 grams per ton ^{*, b}

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ² , not less than.....	_____	%
Salt ² , not more than.....	_____	%
Sodium ³ , not less than	_____	%
Sodium ³ , not more than	_____	%
Potassium, not less than	_____	%
Vitamin A ^{2, 4} , not less than	_____	I.U./lb

¹ When added.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴ Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials (AAFCO).

FEEDING DIRECTIONS

This Type C medicated feed is to be fed continuously as the sole ration for a single, 14 day period at 568 grams to 757 grams tilmicosin per ton on a 100% dry matter basis (511.2 to 681.3 g per ton on a 90% dry matter basis) to provide 12.5 mg tilmicosin/kg of body weight/day and 10 grams to 40 grams monensin per ton on a 90% dry matter basis (11.1 to 44.4 g per ton on a 100% dry matter basis) to provide 0.14 to 0.42 mg monensin/lb bodyweight per day, depending on the severity of the coccidiosis challenge, up to 480 mg/head/day.

CAUTION:

Use only in cattle fed in confinement for slaughter.

Do not allow horses or other equines access to feeds containing tilmicosin or monensin.

Ingestion of monensin by horses has been fatal. Monensin medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. The safety of tilmicosin has not been established in cattle intended for breeding purposes.

Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle.

Must be thoroughly mixed in feeds before use. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

To assure both food safety and responsible use in cattle, the treatment of cattle with this medicated feed is required to be initiated within the first 45 days of the production period. The treatment should not occur concurrent with or following administration of an injectable macrolide, or within 3 days following administration of a non-macrolide injectable BRD therapy

Tilmicosin medicated feed treatment has not been evaluated in cattle with severe clinical disease.

Cattle with severe clinical illness should be evaluated for individual treatment with an alternative non-macrolide therapy.

Complete Type C medicated feeds containing tilmicosin should not be pelleted. Do not use in any feeds containing bentonite, cottonseed meal, or cottonseed hulls. Bentonite, cottonseed meal, or cottonseed hulls in feeds may affect the efficacy of tilmicosin.

The expiration date for a tilmicosin Veterinary Feed Directive (VFD) for cattle must not exceed 45 days from the time of issuance. VFDs for tilmicosin phosphate shall not be refilled.

WARNINGS:

RESIDUE WARNING: Cattle intended for human consumption must not be slaughtered within 28 days of the last treatment with tilmicosin.

Tilmicosin is not approved for use in female dairy cattle 20 months of age or older.

Use in these cattle may cause drug residues in milk. Tilmicosin and monensin are not approved for use in calves intended to be processed for veal. A withdrawal period has not been established in pre-ruminating calves.

Manufactured By
Bluebird Feed Company
Any town, USA 12345

Net Weight: _____ lb (_____ kg)

For emergency medical information, to report an adverse effect, or for technical service call: 1-800-428-4441.

* Final printed label must bear a single drug concentration.

^a 100% dry matter basis.

^b 90% dry matter basis.

^c Sourced from Pulmotil, NADA# 141-064

^d Sourced from Rumensin, NADA# 95-735

Elanco, Pulmotil, Rumensin and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

